

Atlas Progress (updated September 27, 2013)

Block Name	Hours of Coverage (at least 25 hours)	Total Species (observed + breeding birds)	Observed Birds (non-breeding, migrants and winter species)	% Observed Birds (at least 25% non-breeders)	Breeding Birds (at least 50 PO, PR, and CO breeders)	% PR or CO Breeders (at least 67% of all breeding birds)	Comparison to 1985 VA Breeding Bird Atlas (% missing species from Virginia Atlas)
------------	---------------------------------------	---	--	--	--	--	---

KEY:
Red font = area to focus on
 * = Partial block, exceptions may apply

Top Priority Blocks - focus on migrants and winter birds

Ashby Gap - 4*	5.25	48	3	6%	45	36%	22% (29 out of 37 species)
Ashby Gap - 5*	3.7	35	1	3%	34	50%	34% (19 out of 29 species)
Ashby Gap - 6*	42.55	93	10	11%	83	84%	13% (65 out of 75 species)
Charles Town - 6*	11.1	40	1	3%	39	67%	47% (25 out of 47 species)
Round Hill - 2*	28.6	80	12	15%	68	84%	0% (no data from VA Atlas)
Round Hill - 5*	30.35	81	10	12%	71	80%	7% (40 out of 43 species)
Harpers Ferry - 6	39.9	84	15	18%	69	67%	14% (57 out of 66 species)
Purcellville - 1	24.05	83	10	12%	73	74%	2% (42 out of 43 species)
Point of Rocks - 3*	7.9	69	13	19%	56	77%	10% (26 out of 29 species)
Middleburg - 1*	36.05	90	15	17%	75	72%	0% (48 out of 48 species)
Middleburg - 3*	5.5	61	0	0%	61	77%	13% (48 out of 55 species)
Middleburg - 4*	10.2	68	3	4%	65	63%	7% (39 out of 42 species)

needs winter birds

needs winter birds
 needs winter birds

Nearly Complete Blocks - please enter any outstanding data

Bluemont - 2	93.4	99	21	21%	78	73%	12% (14 out of 16 species)
Middleburg - 2	37.2	98	22	22%	76	83%	9% (53 out of 58 species)
Waterford - 6*	41.2	116	45	39%	71	58%	13% (53 out of 61 species)
Leesburg - 1	49.23	107	40	37%	67	58%	11% (47 out of 53 species)
Leesburg - 3	41.85	106	32	30%	74	61%	7% (41 out of 44 species)
Leesburg - 4	37.3	92	23	25%	69	58%	12% (52 out of 59 species)
Sterling - 5	76.79	114	53	46%	61	62%	0% (no VA comparison)

Completed Blocks

Upperville - 2*	29.5	77	23	30%	54	52%	30% (38 out of 54 species)
Bluemont - 1*	161.3	142	59	42%	83	63%	6% (47 out of 50 species)
Bluemont - 3	92.35	131	41	31%	90	84%	7% (64 out of 69)
Bluemont - 4	124.9	148	64	43%	84	90%	4% (50 out of 52 species)
Bluemont - 5	57.65	109	26	24%	83	83%	9% (64 out of 70 species)
Bluemont - 6	48.8	138	49	36%	89	88%	3% (58 out of 60 species)
Rectortown - 1*	20.6	80	21	26%	59	49%	23% (47 out of 61 species)
Rectortown - 2*	23.5	95	21	22%	74	86%	6% (48 out of 51 species)
Round Hill - 4*	116.35	127	46	36%	81	73%	6% (50 out of 53 species)
Round Hill - 6	51.5	105	29	28%	76	80%	12% (56 out of 64 species)
Harpers Ferry - 3*	151.2	130	47	36%	83	89%	11% (50 out of 56 species)
Harpers Ferry - 4*	23.6	95	22	23%	73	73%	6% (30 out of 32 species)
Harpers Ferry - 5	129.2	143	58	41%	85	86%	5% (19 out of 20 species)
Purcellville - 2	96.4	109	49	45%	60	78%	20% (41 out of 51 species)
Purcellville - 3	91.7	105	31	30%	74	72%	5% (41 out of 43 species)
Purcellville - 4	191.85	112	44	39%	68	72%	7% (27 out of 29 species)

Purcellville - 5	104.25	94	26	28%	68	71%	13% (40 out of 46 species)
Purcellville - 6	55.7	93	22	24%	71	76%	14% (55 out of 64 species)
Lincoln - 1	139.8	128	47	37%	81	85%	5% (54 out of 57 species)
Lincoln - 2	41.3	108	29	27%	79	77%	10% (44 out of 49 species)
Lincoln - 3	70.55	105	26	25%	79	77%	5% (52 out of 55 species)
Lincoln - 4	56.35	121	44	36%	77	79%	10% (55 out of 61 species)
Lincoln - 5	68.55	110	35	32%	75	87%	4% (53 out of 55 species)
Lincoln - 6	78.98	109	32	29%	77	74%	6% (59 out of 63 species)
Point of Rocks - 5	116.4	133	58	44%	75	79%	7% (43 out of 46 species)
Point of Rocks - 6*	15.45	76	16	21%	60	68%	23% (49 out of 64 species)
Waterford - 1	448.62	123	41	33%	82	71%	9% (51 out of 56 species)
Waterford - 2	38.7	103	33	32%	70	71%	18% (50 out of 61 species)
Waterford - 3	56.7	115	43	37%	72	78%	6% (46 out of 49 species)
Waterford - 4*	52.1	111	30	27%	81	74%	4% (52 out of 54 species)
Waterford - 5	61.77	114	41	36%	73	68%	2% (51 out of 52 species)
Leesburg - 2*	114.45	96	31	32%	65	75%	25% (43 out of 57 species)
Leesburg - 5	299.4	187	96	51%	91	91%	2% (53 out of 54 species)
Leesburg - 6	185.2	146	69	47%	77	94%	8% (58 out of 63 species)
Arcola - 1	22.52	108	26	24%	82	89%	0% (43 out of 43 species)
Arcola - 2	130.15	187	94	50%	93	95%	0% (41 out of 41 species)
Arcola - 3*	14.75	95	30	32%	65	75%	10% (37 out of 41 species)
Arcola - 4	28.98	107	31	29%	76	68%	3% (37 out of 38 species)
Arcola - 5*	10.75	95	25	26%	70	80%	7% (40 out of 43 species)
Arcola - 6*	32	104	32	31%	72	79%	8% (59 out of 64 species)
Gainesville - 2*	6.05	88	18	20%	70	74%	0% (no VA comparison)
Poolesville - 1*	34.73	106	32	30%	74	72%	0% (no data from VA atlas)
Poolesville - 3*	72.68	112	36	32%	76	62%	0% (10 out of 10)
Sterling - 1*	82.5	102	42	41%	60	60%	0% (no VA comparison)
Sterling - 3*	163.96	163	90	55%	73	67%	8% (56 out of 61 species)
Sterling - 4*	279.87	157	78	50%	79	81%	12% (57 out of 65 species)
Sterling - 6*	193.3	120	58	48%	62	82%	26% (49 out of 66 species)
Herndon - 1	59.85	118	55	47%	63	71%	20% (36 out of 45 species)
Herndon - 2*	118.52	102	40	39%	62	81%	20% (41 out of 51 species)
Herndon - 3*	46.67	85	30	35%	55	64%	11% (25 out of 28 species)
Herndon - 4*	10.5	43	28	65%	15	33%	74% (15 out of 58 species)
Herndon - 5*	46.7	82	31	38%	51	69%	21% (37 out of 47 species)
Seneca - 3*	266.07	153	73	48%	80	73%	6% (47 out of 50 species)
Seneca - 5*	59	80	30	38%	50	72%	23% (37 out of 48 species)

Blocks too small to require coverage

Point of Rocks - 4*	0	0	0	0%	0	0%	0%
Buckeystown - 5*	0	0	0	0%	0	0%	0%